



ADAM™ MC Plus & CellT Plus

Most Accurate Fluorescence Cell Counter

PBMCs | 10 sec per test

Stem cells

15 µL sample loading volume

Primary cells

3.2 µL measuring volume

Cell lines **13 captured images** per channel

ADAM™ MC Plus & CellT Plus are


the highly accurate automated fluorescence cell counter equipped with bright field and two fluorescent channels (AO/DAPI).

4 tests on 1 slide




ADAM™ MC Plus is used for R&D, ADAM™ CellT Plus is available in cGMP production environment.

Bright field & Dual fluorescence (AO/DAPI)




ADAM™ MC Plus & CellT Plus

ADAM™ MC Plus & CellT Plus measures the number of total cells, viable cells, non-viable cells and shows viability results. In addition, they analyzes the cell size and cell aggregation ratio as well.




▲ Cell Size



▲ Cell aggregation ratio

Applicable to a various cell lines


It is possible to use ADAM™ MC Plus & CellT Plus depending on the cell types (PBMCs, etc.) that needs to be monitored during the manufacturing of cell therapy products.



Various applications

Large measuring volume

13 images 3.2 μL measuring volume




Large measurement volume obtained by detecting multiple images of the samples through moving stage provides more accurate results compared to other manufacturers' products.


- · Cell counting accuracy depends on the counting volume (measurement volume).
- The obtained multi-images are processed by image analysis software integrated inside the system.

Accuracy & Repeatability

Correlation of PBMCs total counting between flow cytometry, manual count and $ADAM^{TM}$ MC Plus & CellT Plus.




Data were compared between flow cytometer and ADAM™ MC Plus & CellT Plus results in 20 different PBMCs concentration samples.



Data were compared between manual count and ADAMTM MC Plus results in 20 different PBMCs concentration samples.

Linearity & Reproducibility



A high-concentration (1.8E+07cells/mL) of PBMCs was diluted and a dilution series was counted by ADAM™ MC Plus & CellT Plus. It shows excellent dilution linearity and reproducibility.

	High	Medium	Low
Mean	1.8.E+07	9.0.E+06	1.6.E+06
SD	2.7.E+05	2.3.E+05	3.8.E+04
CV (%)	1.5 %	2.5 %	2.3 %

Sample with low, medium and high concentration of PBMCs were counted with three ADAM™ MC Plus & CellT Plus.

Specifications I





ADAM™ MC Plus & CellT Plus

Hardware		
Measuring range	$5\times10^4 \sim 2\times10^7$ cells/mL	
Optimal range	$4\times10^5 \sim 1\times10^7$ cells/mL	
Analysis time	fast mode 10 sec / test real cell size mode 30 sec / test	
Measuring volume	3.2 μL	
Focus	Auto-focusing	
Objective lens	4 X	
Weight	7.0 kg	
Dimensions (W × D × H)	277 x 276 x 270 mm	

AccuPlus Slide & Reagent

Performance			
Staining method	Acridine orange (AO) & 4',6-diamidino-2-phenylindole (DAPI) stain		
Sample loading volume	15 μL/test		

Ordering Information

Cat. No.	Description	Contents
ADAM-MC Plus	Fluorescence cell analyzer	Main deviceUser manual
ADAM-CellT Plus	Fluorescence cell analyzer for cGMP	 Main device User manual 21 CFR PART 11 requirement support appendix
APAD-400	Cell viability reagent	• 1 mL x 6 tubes (400 Tests)
AP4S-100	AccuPlus Slide 4ch.	4ch. Slide 100 ea



FOR RESEARCH USE ONLY. This product is not approved for diagnostic or therapeutic use.

